

Josephson effect and circle map

Christiansen, Peder Voetmann

Publication date:
1987

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Christiansen, P. V. (1987). *Josephson effect and circle map*. Roskilde Universitet.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

TEKST NR 138

1987

JOSEPHSON EFFECT AND CIRCLE MAP.

Paper presented at
The International Workshop on Teaching Nonlinear Phenomena
at Universities and Schools, "Chaos in Education".
Balaton, Hungary, 26 April - 2 May 1987.

By
Peder Voetmann Christiansen,
Institute of Mathematics and Physics,
Roskilde University Center,
P.O. box 260, DK 4000 Roskilde, Denmark.

TEKSTER fra

IMFUFA **ROSKILDE UNIVERSITETSCENTER**
INSTITUT FOR STUDIET AF MATEMATIK OG FYSIK SAMT DERES
FUNKTIONER I UNDERVISNING, FORSKNING OG ANVENDELSER

JOSEPHSON EFFECT AND CIRCLE MAP.

Peder Voetmann Christiansen

IMFUFA Text no. 138/87

27 pages

ISSN 0106-6242

ABSTRACT.

The Josephson effects associated with a thin insulating layer or a narrow constriction between two superconductors has important technological applications and also provides an instructive example of non-linear effects that may be studied with relatively simple methods. The system is dynamically analogous to a damped pendulum driven by a constant and an oscillating torque and may be simulated by solving a non-linear second order differential equation. The phenomena of phase-locking and transition to chaos are, however, much more conveniently studied by the discrete circle-map.

This paper discusses the connection between the continuous time Josephson equation and the discrete circle-map by the method of return-maps. It will be shown that the so called critical circle map in many cases (but not all) is associated with the transition to hysteretic behaviour of the Josephson junction. The critical circle map, exhibiting a fractal mode-locking structure, is an example of the relevance of the mathematical theories of rational numbers to physics and is recommended as a good example of the challenges of experimental mathematics.

1. The Josephson equation

Superconductivity is one striking example of a macroscopic quantum phenomenon. A crude explanation for metallic superconductivity is obtained by the observation by Cooper (1956) that the effective interaction between electrons through lattice vibrations (phonons) is attractive for states near the Fermi surface, so it becomes energetically favorable to form pair-states, the so called Cooper pairs. These pairs have spin zero and act therefore like particles obeying Bose-Einstein statistics, like e.g. the atoms of He^4 , and similarly to liquid Helium the electrons will condense in a superfluid state at sufficiently low temperatures. This is an example of a Bose-condensation due to the "social" nature of Bose-particles who, quite unlike the "exclusive" Fermi particles, prefer to occupy the same quantum state.

The common wave function for the condensed Cooper-pairs thus acquires the status of a macroscopic order-parameter for the superconducting state, and all of the most striking phenomena of superconductivity, like perfect diamagnetism (Meissner effect), flux quantization, and persistent currents, can be related to the quantum mechanical nature of this complex, scalar field that is coupled to the electromagnetic vector-field. The combined thermodynamic and quantum-mechanical theory leading to non-linear partial differential equations for these two fields was discovered already in 1950 by Landau and Ginzburg¹, and this phenomenological theory has later been able to account for the Josephson effects².

In 1962 it was suggested by Brian D. Josephson that a thin insulating layer between two bulk superconductors should be able to carry a tunnelling supercurrent in the absence of any voltage. The magnitude of this current is given by the dc-equation

$$I_s = I_c \cdot \sin\phi \quad (1)$$

where I_c is the critical current and ϕ is the phase difference between the superconducting wave functions on the two sides of the barrier. In the presence of a voltage difference, V , over the barrier, the phase difference will increase linearly with time in accordance with the ac-equation

$$d\phi/dt = 2eV/\hbar \quad (2)$$

where e is the numerical charge of the electron (thus, the charge of a Cooper-pair is $-2e$) and $\hbar = h/2\pi$, h being Planck's quantum of action. Eqs (1) and (2) together show that a constant dc-voltage over the junction will produce an ac-supercurrent with vanishing mean value.

In practice, however, it is not easy to control the voltage across the junction. The experimental situation is one of current-control, because the resistance of the junction is much smaller than the resistance in the external wires. The junction-resistance is due to the normal (i.e. not condensed) electrons, and this normal fluid is thus providing an extra channel for the current, parallel to the super-channel described by eq. (1). The normal current is given by Ohm's law:

$$I_n = V/R_n \quad (3)$$

where R_n is the normal resistance. Finally, if the junction has a non-vanishing capacitance, C , the following displacement-current must be added to the super- and the normal current:

$$I_d = C \cdot dV/dt \quad (4)$$

The resulting model, called the resistively and capacitively shunted Josephson junction, is very successful in explaining the observed current-voltage characteristics, when a Josephson junction is exposed to a microwave field. The effect of the microwaves is then described by an ac-current source, $I_{AC} \cdot \sin \omega t$, added to the dc-current, I_{DC} . The model in fig. 1a corresponds to the equation:

$$I_d + I_n + I_s = I_{DC} + I_{AC} \cdot \sin \omega t \quad (5)$$

This equation can be reduced to a dimensionless form in the following way. First, we measure the currents in units of the critical current, I_c , putting

$$I = I_{DC}/I_c ; \quad A = I_{AC}/I_c \quad (6)$$

Next, we note that the super-channel, described by eq.s (1) and (2), for small values of the phase ϕ acts like a self-inductance

$$L_s = \hbar/(2eI_c), \quad (7)$$

so the ratio between this quantity and the normal resistance, $\tau = L_s/R_n$, can be used as a unit of time. Thus,

we introduce the symbols

$$\dot{\phi} = \tau \cdot d\phi/dt \quad ; \quad \ddot{\phi} = \tau^2 \cdot d^2\phi/dt^2 \quad ; \quad f = \tau \cdot \omega/2\pi \quad (8)$$

and get the following second order differential equation:

$$Q^2 \cdot \ddot{\phi} + \dot{\phi} + \sin\phi = I + A \cdot \sin(2\pi fT) \quad (9)$$

where $T = t/\tau$, and Q is the dimensionless quantity

$$Q = R_n \cdot \sqrt{2eI_C C/\hbar} \quad (10)$$

As shown in fig. 1b the Josephson equation (9) may just as well describe a classical mechanical system: a damped and driven pendulum. The device may be constructed as metal disc with an excentric mass, M , and a friction unit, F , (e.g. a magnet). Through a torque-control unit, T , the experimenter should be able to apply a torque, consisting of a constant part superposed on a sinusoidally oscillating part, to the pendulum. The correspondence between the parameters and dynamic variables of the two analogous systems is shown in the table below.

<u>Josephson junction</u>	<u>Pendulum</u>
Phase ϕ	Angle from vertical
Supercurrent $I_s = I_C \sin\phi$	Torque from gravity
Normal current	Torque from friction
Capacitance C	Moment of inertia
Applied current $I_{DC} + I_{AC}$	Applied torque

In the following, however, we shall use "Josephson language", mainly because there is a wealth of experimental data to compare with for many different types of superconducting weak links. Before we go on to discuss such data we may remark the following on the different time scales in question. In addition to the inductive relaxation time τ , used as the time unit in eq. (9), and the period of oscillation τ/f of the ac-current (the microwave field), there are two other characteristic times. One is the capacitive relaxation time

$$R_n \cdot C = Q^2 \cdot \tau \quad (11)$$

and the other is the period of oscillation of the undamped "Josephson plasmon" or the undamped pendulum near the stable equilibrium in the absence of driving torques:

$$2\pi \cdot \sqrt{L_S \cdot C} = 2\pi Q \cdot \tau \quad (12)$$

The parameter Q is the quality factor for the plasmon-resonance; the motion will be oscillatory only for $Q > \frac{1}{2}$. A reason for using Q as the parameter instead of "the damping factor", $G = 1/Q$, is that it becomes easier to consider the case $Q = 0$, which is not a case of "infinite damping", but rather of vanishing capacitance. This case is particularly relevant for the case when the Josephson junction has the shape of a "Dayem-bridge", i.e. a narrow constriction in a thin superconducting film. The model in this case degenerates into "the resistively shunted junction", described by the first order differential equation (9) with $Q = 0$.

The experimental current-voltage characteristic with and without a microwave field present are sketched in fig. 2. (For a more detailed account, see ref. 3, and references therein). The figure applies to the cases $Q < 1/2$; for greater values of Q the curves will exhibit hysteresis. The voltage measured is not the instantaneous value of V , but rather its mean value over a long period of time. A dimensionless measure of the voltage is the mean value of the phase derivative $\dot{\phi}$. This is related to the so called winding number, W , i.e. the average number of full turns of the phase per period of the ac-current (we always assume there is an ac-current with frequency f , even though its amplitude A may be zero):

$$W = \lim_{T \rightarrow \infty} \frac{\phi(T) - \phi(0)}{2\pi T \cdot f} = \frac{\langle \dot{\phi} \rangle}{2\pi f} \quad (13)$$

For $Q=0$, $A=0$ the IV-curve is a hyperbola approaching Ohm's law for the normal metal as I goes to infinity, but for $A>0$ there is a step in the current for every voltage that corresponds to an integral value $p/1$ of the winding number W . These are the so called harmonic steps, and their magnitude can be shown to go like the Bessel function J_p of the amplitude A . The supercurrent ($p=0$) should thus be reduced for $A>0$, but for real junctions there will often be an enhancement instead. This enhancement, the Dayem effect, is one of the few qualitative features that are not revealed by eq. (9) but must be explained by the microscopic theory⁴.

The subharmonic steps (not shown in fig. 2) for $W=p/q$, $q>1$, exist for all rational values of W and will be discussed in the following sections.

2. From continuous to discrete time .

The integration of the Josephson equation (9) is in principle straightforward, but very time consuming on a digital computer. A fast and reliable analog computer is the best tool, but this is very expensive, and the programming is rather intricate due to the presence of the $\sin\phi$. In this paper we discuss an alternative approach: the integration is done on an ordinary digital computer of a type that is common in high-schools and private homes (Amstrad 6128), but the results are presented graphically in a way that makes a transition to a discrete time model possible. Afterwards, the qualitative features of phase locking, bifurcations and chaos are investigated by the discrete model, the circle-map.

The method of integration is a 4th order Runge Kutta with variable step-length. When the four parameters Q , I , A and f have been chosen, the integration leads to a definite trajectory for every initial value of ϕ (in the interval from 0 to 2π) and $\dot{\phi}$. As the system is dissipative, the first part of the solution will be transient and has to be integrated out before the results start to have significance for the I-V characteristic. The determination of the observed voltage will then require a long period of integration before the mean value of $\dot{\phi}$ has attained a reasonably stable value. So a single I-V characteristic, where I is the only parameter to be varied is a tedious job. In the following we keep the ac-amplitude and the driving frequency fixed, $A = 1$, $f = 0.4$.

The period of the ac-current, $1/f$, is the only natural unit of discrete time. At all the times 0 , $1/f$, $2/f$, $3/f$, -

the situation with respect to the external world is the same. If we could find a simple mapping from the state $(\phi, \dot{\phi})$ at one of these times to the state at the next time in the discrete series, the whole investigation could be made much faster.

For the case $Q = 0$ the situation is even simpler, because ϕ is the only state variable. Instead of ϕ we use the reduced phase

$$y_n = \phi(n/f)/2\pi \quad (14)$$

so when ϕ is confined to the interval from 0 to 2π , y will lie between 0 and 1. The return map, F , defined by

$$y_{n+1} = F(y_n) \quad (15)$$

will then be well defined, and because there is a unique solution curve for every ϕ at a given time, we can be sure that F is a monotonous function. In some cases the discrete solutions y_n will repeat themselves after a small period and a graphical representation of y_{n+1} versus y_n will only show isolated dots, but in other cases a whole curve can be traced out. Fig. 3 shows such a case.

In general the return map for the Josephson equation, when it exists, can be shown to be a circle map⁵, i.e. a non-linear mapping of the unit circle on itself. The function F has two points of inflection and is reasonably well described as a case of Arnold's sine map⁶:

$$y_{n+1} = y_n + \Omega - (K/2\pi) \cdot \sin(2\pi \cdot y_n) \quad (16)$$

where the values of the dynamical variable y have to be reduced to the interval between 0 and 1. When comparing the return maps (15) to the sine map (16) one must allow for an arbitrariness in the choice of the phase of the ac-current at the discrete instants picked out. This means that there will be an arbitrary offset in the y -scale, so that the two points of inflection do not occur for $y=0$ and $y=1/2$, as they do for the sine map. This is, however, not essential.

The surprising thing is that the sine map fits the return maps so well, even when $Q > 0$ and the state space is two dimensional. The explanation is that the dissipative character of the system ensures that the discrete motion in the $(\phi, \dot{\phi})$ plane after the transient period will be restricted to an attractor of dimension lower than 2, and that in typical cases this attractor will define $\dot{\phi}$ as a single-valued function of ϕ . For $Q=0$ we have according to (9):

$$\dot{\phi}_n = I - \sin \phi_n \quad (17)$$

so for small values of Q we can expect something similar.

For $Q=0$ the return map is necessarily increasing, so the parameter K for the approximating sine map is less than 1. The circle map is then subcritical. The value of K is one minus the slope of the tangent at the point of inflection (comp. fig. 3). For $Q>0$ we may find a supercritical return map with $K>1$ and negative slope of the tangent, as shown in fig. 4. Correspondingly, in such cases the discrete attractor in the $(\phi, \dot{\phi})$ plane for some values of ϕ allows more than one value of $\dot{\phi}$, as can be seen in fig. 5. A closer inspection of the supercritical return

maps reveal small wiggles, so they are not strictly univalued⁵, but for a slightly supercritical case the wiggles are too small to be seen on a normal scale and are not important for the gross features of the I-V characteristics. We conclude that the return maps for the Josephson equation are well described by the sine map for the subcritical, critical and slightly supercritical cases.

When investigating the dynamics of the sine map (16) one normally chooses a fixed value of K and determines the winding number

$$W = \lim_{n \rightarrow \infty} (y_n - y_0)/n \quad (18)$$

as a function of the parameter Ω . It then turns out that phase-locking occurs for every rational value $W = p/q$ in a finite Ω -interval. These phase locking intervals correspond to the steps on the Josephson I-V characteristics, both harmonic and subharmonic, but the correspondence is not straightforward, because when we vary I alone, both Ω and K change; $\Omega(I)$ is increasing, while $K(I)$ is decreasing. For some values of Q a transition takes place from supercritical to subcritical behaviour when I increases beyond a value $I_1(Q)$. The transition can only be seen on the I-V curves when $I_1(Q)$ is greater than the smallest current that allows a solution with nonzero voltage. For $A=1$, $f=0.4$ this will be the case in the interval $0.4 < Q < 1.1$; I_1 has a maximum ≈ 1.3 for $Q \approx 2$. No supercritical parts can be found on the I-V characteristics for these values of A and f with winding numbers above 0.44.

Fig. 6 shows three different return maps for the same

current $I = 1.05$ but three different Q -values, 0.3, 1, and 1.5. For $Q = 1$ we see only three isolated groups of points of three points each. This is a case of phase-locking and the winding number is $1/3$, but as there are nine points the mode structure is not $1/3$, but $3/9$. For a supercritical case there are several modes for some of the winding numbers that correspond to irreducible fractions p/q . This is because there is a local maximum on the circle map for $K > 1$, and therefore we can expect to find traces of the Feigenbaum route to chaos superposed on the mode locked steps.

For the subcritical circle map there is a finite measure of Ω -values giving an irrational winding number, but in the critical case, $K = 1$, this measure is zero and the values form a Cantor-set of fractal dimension 0.87, as will be discussed in the next section. For supercritical cases some of the phase-locked Ω -intervals overlap, meaning that we can find different winding numbers for the same Ω . This gives rise to hysteresis in the Josephson I - V curves, a feature that can be used for distinguishing supercritical regions from subcritical ones when looking at experimental data. However, there is another sort of hysteresis as regards the supercurrent for $Q > 1$. For $Q = 1.5$ we can lower the current from above 1 down to about 0.7 and find a subcritical circle map all the way, until it touches the line $y_{n+1} = y_n$ at which point the transition to zero voltage suddenly takes place. Raising the current again we stay on the supercurrent (the 0/1 step) until the current is almost 1, when the winding number changes discontinuously from 0 to 0.41. Apparently the return map between these two transitions includes an isolated point on the line.

3. The critical sine map .

The winding number W as a function of the driving parameter α for the sine map forms a "Devil's staircase". For the critical case $K = 1$ the staircase is complete, which means that the steps for rational W together have the same measure as the α -domain, although they do not cover it completely but leave a Cantor-set of zero measure. Considering α as a function of W we have the curious case of a function that is discontinuous for all rational values, but continuous for all irrational values of W . Fig. 7 shows a graph of the complete staircase for $0 < \alpha < \frac{1}{2}$. This is sufficient, because $W(1-\alpha) = 1 - W(\alpha)$.

The dynamical process for the critical sine map can be depicted as in fig. 8, where subsequent values of y (reduced to the interval between 0 and 1) have been plotted while α slowly increases from 0 to $\frac{1}{2}$. There is no real "chaos" in this picture, although it may look so. An irrational winding number corresponds to a quite regular motion on the circle, although it never repeats itself.

The critical sine map is particularly interesting because it shows a universal scaling structure common to all circle maps with a horizontal slope at the point of inflection, provided that the local behaviour is cubic. Shenker showed⁷ for the Fibonacci-fractions q_n/q_{n+1} , where the numerator and the denominator are consecutive members of the Fibonacci-sequence

$$1, 1, 2, 3, 5, 8, \dots, q_n = q_{n-1} + q_{n-2}, \dots \quad (19)$$

that the stability intervals asymptotically scale with the

power law

$$\Delta \Omega(q_{n-1}/q_n) \propto q_n^{-2.164} \quad (20)$$

If all stability intervals for $W = p/q$ scaled with the denominator in the same power, $-\alpha$, we could easily calculate the fractal dimension for the set of qs with irrational W . If $L(\Delta)$ denotes the measure seen when all stability intervals less than Δ are ignored, we have

$$dL(\Delta)/d\Delta \propto \tilde{q}(\Delta)^{1-\alpha} \cdot d\tilde{q}(\Delta)/d\Delta \quad (21)$$

where we have used that the number of irreducible fractions with denominator q is proportional to q , asymptotically. The quantity $\tilde{q}(\Delta)$ is the largest denominator whose stability interval can be seen with the Δ -scale, i.e. $\tilde{q} \propto \Delta^{-1/\alpha}$, and therefore

$$dL(\Delta)/d\Delta \propto \Delta^{-2/\alpha} \quad (22)$$

But by definition⁸ of D , the fractal dimension, $L \propto \Delta^{1-D}$, so we should have

$$D = 2/\alpha \quad (23)$$

Shenker's value, $\alpha = 2.164$ would thus give $D = 0.924$. A direct measurement using calculated stability intervals and the definition of D gives however⁹,

$$D = 0.870 \quad (24)$$

so, it seems that the Fibonacci fractions studied by Shenker are not very "typical". In fact they are extreme in the

sense that the Fibonacci intervals are always the largest for fractions with the same denominator. Another extreme are the fractions $1/q$ which always have the smallest intervals for the given q . These intervals are easily shown to go like q^{-3} , a consequence of the fact that N rises like a square root on the lowest part of the Devil's staircase.

The best way to define what is meant by "extreme" or "typical" fractions is based on the continued fraction expansion

$$p/q = 1/(a_1 + 1/(a_2 + 1/(a_3 + \dots + 1/a_N))) \dots \quad (25)$$

The number of terms, $N(p/q)$ can be shown to lie between the limits 1 and $N_{\max}(q)$, where $N_{\max}(q)$ is the number characterizing the Fibonacci-fractions for which all the terms are 1. Asymptotically for q large:

$$N_{\max}(q) = \frac{\ln[(3-\tau) \cdot q]}{\ln \tau} \quad (26)$$

where $\tau = (\sqrt{5}+1)/2$ is the golden mean ($1/\tau$ is the limiting value of the Fibonacci-fractions). An even better way of characterizing the fractions is to use the sum of terms

$$S = \sum a_n \quad (27)$$

For given q , S will lie between the maximum q (for $1/q$) and the minimum, attained by the Fibonacci-fractions

$$S_{\min}(q) = N_{\max}(q) \quad (28)$$

As a rule of thumb one can say that the length of the

stability intervals $\Delta\Omega(p/q)$ depends only on q and S , i.e. if two fractions have identical values of q and S we can be sure that the corresponding stability intervals have equal lengths within a few percent, even if their N values differ.

A way of characterizing "typical" fraction is suggested by the result¹⁰ that the values of N for large q are normally distributed around the value

$$N_t(q) = \frac{12 \ln 2}{\pi^2} \cdot \ln q \quad (29)$$

with a standard deviation given by $\ln N_t$. The value N_t is about 40% of N_{\max} . If we look at the quantity

$$\sigma(q, S) = [1 + (S - S_{\min}) / (N_t \ln N_t)]^{-1} \quad (30)$$

we can see that it is confined to the interval between 0 and 1. Furthermore, it will have a rather stable distribution around the "typical" value $\frac{1}{2}$. An empirical formula for the phase-locking intervals is given by

$$\Delta\Omega(p/q) \approx b \cdot (q/r)^{-\delta(\sigma)} \quad (31)$$

where $b \approx 0.035$ and $r \approx 2.8$. The exponent δ is given by the expression (30) using the interpolation formula

$$\delta(\sigma) \approx 2 + 1 / (5.1 \cdot \sigma + 1) \quad (32)$$

The expressions (30)-(32) summarize some of the main known features of the critical sine map and give within 10% uncertainties the correct magnitudes of the intervals, at least up to denominators about 100. However, it does not

pretend to give any exact insight in the self-similar structure. The self-similarity is not strict, but slightly modulated from place to place on the staircase, but still, the overall fractal dimension $D = 0.870$ seems very well defined. The universality of this result is confirmed by direct measurements of D on the critical steps of IV-characteristics for Josephson junctions simulated on an analog computer¹¹.

A proper theory for the scaling properties may lie hidden in the particular ordering of rational numbers known as the Farey-tree¹². For a given value of the sum S ($S > 1$) there are exactly 2^{S-2} irreducible fractions between 0 and 1, which suggests that all these fractions may be ordered in a binary tree where each layer has a constant S . As shown in fig. 9 the construction is very simple: a fraction is formed by adding both the numerators and the denominators of the nearest adjacent fractions on the left and the right in the layers above. This will preserve the natural ordering of all the numbers from left to right. Fig. 9 shows also how to determine the number of terms in the continued fraction expansion by visual inspection: The branches of the tree are drawn alternating between solid and dashed lines. By starting in the point $1/1$ where $N=1$ and going to another point p/q , N is increased by one for every solid line and stays constant on the dashed lines.

The number theoretical mysteries hidden in such hard boiled physical problems as the Josephson junction and the pendulum have just begun to reveal themselves through the experimental mathematics of the circle maps.

Acknowledgements .


The author gratefully acknowledges stimulating discussions with T. Bohr, P. Cvitanovic, and M. T. Levinsen.


References .

1. L. D. Landau and V. L. Ginzburg, Zh. Eksperim. i Teor. Fiz., 20 , 1064 (1950).
2. P. V. Christiansen, E. B. Hansen and C. J. Sjöström, J. Low Temp. Phys., 4 , 349 (1971).
3. D. E. McCumber, J. Appl. Phys., 39 , 3113 (1968).
4. G. M. Eliashberg, Zh. Eksperim. i Teor. Fiz. Pis. Red., 11 , 186 (1970).
5. P. Bak, T. Bohr, M. H. Jensen and P. V. Christiansen, Solid State Comm., 51 , 231 (1984).
6. V. I. Arnold, Izv. Acad. Nauk., 25 , 21 (1961); Trans. Am. Math. Soc., 46 , 213 (1965).
7. S. J. Shenker, Physica, 5D , 405 (1982).
8. B. Mandelbrot, "Fractals: Form, Chance, and Dimension", W. H. Freeman, San Fransisco (1977).
9. M. H. Jensen, P. Bak and T. Bohr, Phys. Rev. Lett., 50 , 1637 (1983).
10. A. I. Khinchin, "Kettenbrüche", B. G. Teubner, Leipzig (1956).
11. W. J. Yeh, D. a-Ren He and Y. H. Kao, Phys. Rev. Lett., 52 , 480 (1984).
P. Alstrøm and M. T. Levinsen, Phys. Rev. B 31 , 2753 (1985); *ibid.* 32 , 1503 (1985).
12. P. Cvitanovic and B. Shraiman, private communication.

Figure captions .


- Fig. 1 . Iconic models of physical analogies for the Josephson equation.
- Fig. 2 . Current (I) - voltage (V) characteristics for Josephson junction. Solid curve: without ac-current. Dotted curve: with ac-current. Line: normal conductivity.
- Fig. 3 . Return map for $Q=0$ showing how to determine K by the slope of the tangent at the point of inflection.
- Fig. 4 . Supercritical return map.
- Fig. 5 . Discrete attractor corresponding to the return map of fig. 4. Horizontal: ϕ , vertical: $\dot{\phi}$. The arrow indicates a place where $\dot{\phi}$ is not uniquely determined from ϕ .
- Fig. 6 . Three return maps for the same current.
- Fig. 7 . Devil's staircase for the critical sine map.
- Fig. 8 . Dynamics of the critical sine map.
- Fig. 9 . The Farey-tree.






(up)

Fig. 2




(up)

Fig. 3




(up)

Fig. 4




(up)

Fig. 5




(up)

Fig. 6




(up)

Fig. 7



(up)

Fig. 8



(up)

Fig. 9

- 1/78 "TANKER OM EN PRAKSIS" - et matematikprojekt.
Projektrapport af: Anne Jensen, Lena Lindenskov, Marianne Kesselhahn og Nicolai Lomholt.
Vejleder: Anders Madsen
- 2/78 "OPTIMERING" - Menneskets forøgede beherskelsesmuligheder af natur og samfund.
Projektrapport af: Tom J. Andersen, Tommy R. Andersen, Gert Krenø og Peter H. Lassen
Vejleder: Bernhelm Boss.
- 3/78 "OPCAVESAMLING", breddekursus i fysik.
Af: Lasse Rasmussen, Aage Bonde Kræmmer og Jens Højgaard Jensen.
- 4/78 "TRE ESSAYS" - om matematikundervisning, matematiklæreruddannelsen og videnskabsrindalismen.
Af: Mogens Niss
Nr. 4 er p.t. udgået.
- 5/78 "BIBLIOGRAFISK VEJLEDNING til studiet af DEN MODERNE FYSIKS HISTORIE".
Af: Helge Kragh.
Nr. 5 er p.t. udgået.
- 6/78 "NOGLE ARTIKLER OG DEBATINDLÆG OM - læreruddannelse og undervisning i fysik, og - de naturvidenskabelige fags situation efter studenteroprøret".
Af: Karin Beyer, Jens Højgaard Jensen og Bent C. Jørgensen.
- 7/78 "MATEMATIKKENS FORHOLD TIL SAMFUNDSØKONOMIEN".
Af: B.V. Gnedenko.
Nr. 7 er udgået.
- 8/78 "DYNAMIK OG DIAGRAMMER". Introduktion til energy-bond-graph formalismen.
Af: Peder Voetmann Christiansen.
- 9/78 "OM PRAKSIS' INDFLYDELSE PÅ MATEMATIKKENS UDVIKLING". - Motiver til Kepler's: "Nova Stereometria Doliorum Vinariorum".
Projektrapport af: Lasse Rasmussen.
Vejleder: Anders Madsen.
-
- 10/79 "THERMODYNAMIK I GYMNASIET".
Projektrapport af: Jan Christensen og Jeanne Mortensen.
Vejledere: Karin Beyer og Peder Voetmann Christiansen.
- 11/79 "STATISTISKE MATERIALER".
Af: Jørgen Larsen.
- 12/79 "LINEÆRE DIFFERENTIALLIGNINGER OG DIFFERENTIALLIGNINGSSYSTEMER".
Af: Mogens Brun Heefelt.
Nr. 12 er udgået.
- 13/79 "CAVENDISH'S FORSØG I GYMNASIET".
Projektrapport af: Gert Kreinø.
Vejleder: Albert Chr. Paulsen.
- 14/79 "BOOKS ABOUT MATHEMATICS: History, Philosophy, Education, Models, System Theory, and Works of".
Af: Else Høyrup.
Nr. 14 er p.t. udgået.
- 15/79 "STRUKTUREL STABILITET OG KATASTROFER i systemer i og udenfor termodynamisk ligevægt".
Specialeopgave af: Leif S. Striegler.
Vejleder: Peder Voetmann Christiansen.
- 16/79 "STATISTIK I KRÆFTFORSKNINGEN".
Projektrapport af: Michael Olsen og Jørn Jensen.
Vejleder: Jørgen Larsen.
- 17/79 "AT SPØRGE OG AT SVARE i fysikundervisningen".
Af: Albert Christian Paulsen.
- 18/79 "MATHEMATICS AND THE REAL WORLD", Proceedings af an International Workshop, Roskilde University Centre, Denmark, 1978.
Preprint.
Af: Bernhelm Booss og Mogens Niss (eds.)
- 19/79 "GEOMETRI, SKOLE OG VIRKELIGHED".
Projektrapport af: Tom J. Andersen, Tommy R. Andersen og Per H.H. Larsen.
Vejleder: Mogens Niss.
- 20/79 "STATISTISKE MODELLER TIL BESTEMMELSE AF SIKRE DOSER FOR CARCINOGENE STOFFER".
Projektrapport af: Michael Olsen og Jørn Jensen.
Vejleder: Jørgen Larsen
- 21/79 "KONTROL I GYMNASIET-FORMÅL OG KONSEKVENSER".
Projektrapport af: Crilles Bacher, Per S. Jensen, Preben Jensen og Torben Nysteen.
- 22/79 "SEMIOTIK OG SYSTEMEGENSKABER (1)".
1-port lineært response og støj i fysikken.
Af: Peder Voetmann Christiansen.
- 23/79 "ON THE HISTORY OF EARLY WAVE MECHANICS - with special emphasis on the role of reality".
Af: Helge Kragh.
-
- 24/80 "MATEMATIKOPFATTELSE HOS 2. G'ERE".
a+b 1. En analyse. 2. Interviewmateriale.
Projektrapport af: Jan Christensen og Knud Lindhardt Rasmussen.
Vejleder: Mogens Niss.
- 25/80 "EKSAMENSOPGAVER", Dybdemodulet/fysik 1974-79.
- 26/80 "OM MATEMATISKE MODELLER".
En projektrapport og to artikler.
Af: Jens Højgaard Jensen m.fl.
- 27/80 "METHODOLOGY AND PHILOSOPHY OF SCIENCE IN PAUL DIRAC'S PHYSICS".
Af: Helge Kragh.
- 28/80 "DILENTRISK RELAXATION - et forslag til en ny model bygget på væskernes viscoelastiske egenskaber".
Projektrapport af: Gert Kreinø.
Vejleder: Niels Boye Olsen.
- 29/80 "ODIN - undervisningsmateriale til et kursus i differentiaalligningsmodeller".
Projektrapport af: Tommy R. Andersen, Per H.H. Larsen og Peter H. Lassen.
Vejleder: Mogens Brun Heefelt.
- 30/80 "FUSIONSENERGIEN - - - ATOMSAMFUNDETS ENDESTATION".
Af: Oluf Danielsen.
Nr. 30 er udgået.
- 31/80 "VIDENSKABSTEORETISKE PROBLEMER VED UNDERVISNINGSSYSTEMER BASERET PÅ MÆNGDELÆRE".
Projektrapport af: Troels Lange og Jørgen Karrebæk.
Vejleder: Stig Andur Pedersen.
Nr. 31 er p.t. udgået.
- 32/80 "POLYMERE STOFFERS VISCOELASTISKE EGENSKABER - BELYST VED HJÆLP AF MEKANISKE IMPEDANSMÅLINGER - MØSSBAUEREFFEKTMÅLINGER".
Projektrapport af: Crilles Bacher og Preben Jensen.
Vejledere: Niels Boye Olsen og Peder Voetmann Christiansen.
- 33/80 "KONSTITUERING AF FAG INDEN FOR TEKNISK - NATURVIDENSKABELIGE UDDANNELSER. I-II".
Af: Arne Jakobsen.
- 34/80 "ENVIRONMENTAL IMPACT OF WIND ENERGY UTILIZATION".
ENERGY SERIES NO. I.
Af: Bent Sørensen
Nr. 34 er udgået.

- 35/80 "HISTORISKE STUDIER I DEN NYERE ATOMFYSIKS UDVIKLING".
Af: Helge Kragh.
- 36/80 "HVAD ER MENINGEN MED MATEMATIKUNDERVISNINGEN?".
Fire artikler.
Af: Mogens Niss.
- 37/80 "RENEWABLE ENERGY AND ENERGY STORAGE".
ENERGY SERIES NO. 2.
Af: Bent Sørensen.
-
- 38/81 "TIL EN HISTORIE TEORI OM NATURERKENDELSE, TEKNOLOGI OG SAMFUND".
Projektrapport af: Erik Gade, Hans Hedal, Henrik Lau og Finn Physant.
Vejledere: Stig Andur Pedersen, Helge Kragh og Ib Thiersen.
Nr. 38 er p.t. udgået.
- 39/81 "TIL KRITIKKEN AF VÆKSTØKONOMIEN".
Af: Jens Højgaard Jensen.
- 40/81 "TELEKOMMUNIKATION I DANMARK - oplæg til en teknologivurdering".
Projektrapport af: Arne Jørgensen, Bruno Petersen og Jan Vedde.
Vejleder: Per Nørgaard.
- 41/81 "PLANNING AND POLICY CONSIDERATIONS RELATED TO THE INTRODUCTION OF RENEWABLE ENERGY SOURCES INTO ENERGY SUPPLY SYSTEMS".
ENERGY SERIES NO. 3.
Af: Bent Sørensen.
- 42/81 "VIDENSKAB TEORI SAMFUND - En introduktion til materialistiske videnskabsopfattelser".
Af: Helge Kragh og Stig Andur Pedersen.
- 43/81 1. "COMPARATIVE RISK ASSESSMENT OF TOTAL ENERGY SYSTEMS".
2. "ADVANTAGES AND DISADVANTAGES OF DECENTRALIZATION".
ENERGY SERIES NO. 4.
Af: Bent Sørensen.
- 44/81 "HISTORISKE UNDERSØGELSER AF DE EKSPERIMENTELLE FORUDSÆTNINGER FOR RUTHERFORDS ATOMMODEL".
Projektrapport af: Niels Thor Nielsen.
Vejleder: Bent C. Jørgensen.
-
- 45/82 Er aldrig udkommet.
- 46/82 "EKSEMPLARISK UNDERVISNING OG FYSISK ERKENDELSE-
1+1 ILLUSTRERET VED TO EKSEMPLER".
Projektrapport af: Torben O. Olsen, Lasse Rasmussen og Niels Dreyer Sørensen.
Vejleder: Bent C. Jørgensen.
- 47/82 "BARSEBÄCK OG DET VÆRST OFFICIELT-TÆNKELIGE UHELD".
ENERGY SERIES NO. 5.
Af: Bent Sørensen.
- 48/82 "EN UNDERSØGELSE AF MATEMATIKUNDERVISNINGEN PÅ ADGANGSKURSUS TIL KØBENHAVNS TEKNISKUM".
Projektrapport af: Lis Eilertzen, Jørgen Karrebæk, Troels Lange, Preben Nørregaard, Lissi Pedersen, Laust Rishøj, Lill Røn og Isac Showiki.
Vejleder: Mogens Niss.
- 49/82 "ANALYSE AF MULTISPEKTRALE SATELLITBILLEDER".
Projektrapport af: Preben Nørregaard.
Vejledere: Jørgen Larsen og Rasmus Ole Rasmussen.
- 50/82 "HERSLEV - MULIGHEDER FOR VEDVARENDE ENERGI I EN LANDSBY".
ENERGY SERIES NO. 6.
Rapport af: Bent Christensen, Bent Hove Jensen, Dennis B. Møller, Bjarne Laursen, Bjarne Lillethorup og Jacob Mørch Pedersen.
Vejleder: Bent Sørensen.
- 51/82 "HVAD KAN DER GØRES FOR AT AFHJÆLPE PIGERS BLOKERING OVERFOR MATEMATIK?".
Projektrapport af: Lis Eilertzen, Lissi Pedersen, Lill Røn og Susanne Stender.
- 52/82 "DESUSPENSION OF SPLITTING ELLIPTIC SYMBOLS".
Af: Bernhelm Booss og Krzysztof Wojciechowski.
- 53/82 "THE CONSTITUTION OF SUBJECTS IN ENGINEERING EDUCATION".
Af: Arne Jacobsen og Stig Andur Pedersen.
- 54/82 "FUTURES RESEARCH" - A Philosophical Analysis of Its Subject-Matter and Methods.
Af: Stig Andur Pedersen og Johannes Witt-Hansen.
- 55/82 "MATEMATISKE MODELLER" - Litteratur på Roskilde Universitetsbibliotek.
En biografi.
Af: Else Højrup.

Vedr. tekst nr. 55/82 se også tekst nr. 62/83.
- 56/82 "EN - TO - MANGE" -
En undersøgelse af matematisk økologi.
Projektrapport af: Troels Lange.
Vejleder: Anders Madsen.
-
- 57/83 "ASPECT EKSPERIMENTET"-
Skjulte variable i kvantemekanikken?
Projektrapport af: Tom Juul Andersen.
Vejleder: Peder Voetmann Christiansen.
Nr. 57 er udgået.
- 58/83 "MATEMATISKE VANDRINGER" - Modelbetragtninger over spredning af dyr mellem småbiotoper i agerlandet.
Projektrapport af: Per Hammershøj Jensen og Lene Vagn Rasmussen.
Vejleder: Jørgen Larsen.
- 59/83 "THE METHODOLOGY OF ENERGY PLANNING".
ENERGY SERIES NO. 7.
Af: Bent Sørensen.
- 60/83 "MATEMATISK MODEKSPERTISE"- et eksempel.
Projektrapport af: Erik O. Gade, Jørgen Karrebæk og Preben Nørregaard.
Vejleder: Anders Madsen.
- 61/83 "FYSIKS IDEOLOGISKE FUNKTION, SOM ET EKSEMPEL PÅ EN NATURVIDENSKAB - HISTORISK SET".
Projektrapport af: Annette Post Nielsen.
Vejledere: Jens Højrup, Jens Højgaard Jensen og Jørgen Vogelius.
- 62/83 "MATEMATISKE MODELLER" - Litteratur på Roskilde Universitetsbibliotek.
En biografi 2. rev. udgave.
Af: Else Højrup.
- 63/83 "CREATING ENERGY FUTURES: A SHORT GUIDE TO ENERGY PLANNING".
ENERGY SERIES No. 8.
Af: David Crossley og Bent Sørensen.
- 64/83 "VON MATEMATIK UND KRIEG".
Af: Bernhelm Booss og Jens Højrup.
- 65/83 "ANVENDT MATEMATIK - TEORI ELLER PRAKSIS".
Projektrapport af: Per Hedegård Andersen, Kirsten Habekost, Carsten Holst-Jensen, Annelise von Moos, Else Marie Pedersen og Erling Møller Pedersen.
Vejledere: Bernhelm Booss og Klaus Grünbaum.
- 66/83 "MATEMATISKE MODELLER FOR PERIODISK SELEKTION I ESCHERICHIA COLI".
Projektrapport af: Hanne Lisbet Andersen, Ole Richard Jensen og Klavs Frisdahl.
Vejledere: Jørgen Larsen og Anders Hede Madsen.
- 67/83 "ELEPSOIDE METODEN - EN NY METODE TIL LINEÆR PROGRAMMERING?".
Projektrapport af: Lone Billmann og Lars Boye.
Vejleder: Mogens Brun Heefelt.
- 68/83 "STOKASTISKE MODELLER I POPULATIONSGENETIK" - til kritikken af teoriladede modeller.
Projektrapport af: Lise Odgård Gade, Susanne Hansen, Michael Hviid og Frank Mølgård Olsen.
Vejleder: Jørgen Larsen.

- 69/83 "ELEVFORUDSÆTNINGER I FYSIK"
- en test i l.g med kommentarer.
Af: Albert C. Paulsen.
- 70/83 "INDLÆRINGS - OG FORMIDLINGSPROBLEMER I MATEMATIK PÅ VOKSENUNDERVISNINGSNIVEAU".
Projektrapport af: Hanne Lisbet Andersen, Torben J. Andreasen, Svend Åge Houmann, Helle Glerup Jensen, Keld Fl. Nielsen, Lene Vagn Rasmussen.
Vejleder: Klaus Grünbaum og Anders Hede Madsen.
- 71/83 "PIGER OG FYSIK"
- et problem og en udfordring for skolen?
Af: Karin Beyer, Sussanne Blegaa, Birthe Olsen, Jette Reich og Mette Vedelsby.
- 72/83 "VERDEN IFØLGE PEIRCE" - to metafysiske essays, om og af C.S Peirce.
Af: Peder Voetmann Christiansen.
- 73/83 "EN ENERGIANALYSE AF LANDBRUG"
- økologisk contra traditionelt.
ENERGY SERIES NO. 9
Specialeopgave i fysik af: Bent Hove Jensen.
Vejleder: Bent Sørensen.
-
- 74/84 "MINIATURISERING AF MIKROELEKTRONIK" - om videnskabeliggjort teknologi og nytten af at lære fysik.
Projektrapport af: Bodil Harder og Linda Szkotak Jensen.
Vejledere: Jens Højgaard Jensen og Bent C. Jørgensen.
- 75/84 "MATEMATIKUNDERVISNINGEN I FREMTIDENS GYMNASIUM"
- Case: Lineær programmering.
Projektrapport af: Morten Blomhøj, Klavs Frisdahl og Frank Mølgaard Olsen.
Vejledere: Mogens Brun Heefelt og Jens Bjørneboe.
- 76/84 "KERNEKRAFT I DANMARK?" - Et høringsvar indkaldt af miljøministeriet, med kritik af miljøstyrelsens rapporter af 15. marts 1984.
ENERGY SERIES No. 10
Af: Niels Boye Olsen og Bent Sørensen.
- 77/84 "POLITISKE INDEKS - FUP ELLER FAKTA?"
Opinionsundersøgelser belyst ved statistiske modeller.
Projektrapport af: Svend Åge Houmann, Keld Nielsen og Susanne Stender.
Vejledere: Jørgen Larsen og Jens Bjørneboe.
- 78/84 "JÆVNSTRØMSLEDNINGSEVNE OG GITTERSTRUKTUR I AMORFT GERMANIUM".
Specialrapport af: Hans Heddal, Frank C. Ludvigsen og Finn C. Physant.
Vejleder: Niels Boye Olsen.
- 79/84 "MATEMATIK OG ALMENDANNELSE".
Projektrapport af: Henrik Coster, Mikael Wennerberg Johansen, Povl Kattler, Birgitte Lydholm og Morten Overgaard Nielsen.
Vejleder: Bernhelm Booss.
- 80/84 "KURSUSMATERIALE TIL MATEMATIK B".
Af: Mogens Brun Heefelt.
- 81/84 "FREKVENSAFHÆNGIG LEDNINGSEVNE I AMORFT GERMANIUM".
Specialrapport af: Jørgen Wind Petersen og Jan Christensen.
Vejleder: Niels Boye Olsen.
- 82/84 "MATEMATIK - OG FYSIKUNDERVISNINGEN I DET AUTOMATISEREDE SAMFUND".
Rapport fra et seminar afholdt i Hvidovre 25-27 april 1983.
Red.: Jens Højgaard Jensen, Bent C. Jørgensen og Mogens Niss.
- 83/84 "ON THE QUANTIFICATION OF SECURITY":
PEACE RESEARCH SERIES NO. 1
Af: Bent Sørensen
nr. 83 er p.t. udgået
- 84/84 "NOGLE ARTIKLER OM MATEMATIK, FYSIK OG ALMENDANNELSE".
Af: Jens Højgaard Jensen, Mogens Niss m. fl.
- 85/84 "CENTRIFUGALREGULATORER OG MATEMATIK".
Specialrapport af: Per Hødegård Andersen, Carsten Holst-Jensen, Else Marie Pedersen og Erling Møller Pedersen.
Vejleder: Stig Andur Pedersen.
- 86/84 "SECURITY IMPLICATIONS OF ALTERNATIVE DEFENSE OPTIONS FOR WESTERN EUROPE".
PEACE RESEARCH SERIES NO. 2
Af: Bent Sørensen.
- 87/84 "A SIMPLE MODEL OF AC HOPPING CONDUCTIVITY IN DISORDERED SOLIDS".
Af: Jeppe C. Dyre.
- 88/84 "RISE, FALL AND RESURRECTION OF INFINITESIMALS".
Af: Detlef Laugwitz.
- 89/84 "FUERNVARMEOPTIMERING".
Af: Bjarne Lillethorup og Jacob Mørch Pedersen.
- 90/84 "ENERGI I L.G - EN TEORI FOR TILRETTELÆGGELSE".
Af: Albert Chr. Paulsen.
-
- 91/85 "KVANTETEORI FOR GYMNASIET".
1. Lærervejledning
Projektrapport af: Biger Lundgren, Henning Sten Hansen og John Johansson.
Vejleder: Torsten Meyer.
- 92/85 "KVANTETEORI FOR GYMNASIET".
2. Materiale
Projektrapport af: Biger Lundgren, Henning Sten Hansen og John Johansson.
Vejleder: Torsten Meyer.
- 93/85 "THE SEMIOTICS OF QUANTUM - NON - LOCALITY".
Af: Peder Voetmann Christiansen.
- 94/85 "TREENIGHEDEN BOURBAKI - generalen, matematikeren og ånden".
Projektrapport af: Morten Blomhøj, Klavs Frisdahl og Frank M. Olsen.
Vejleder: Mogens Niss.
- 95/85 "AN ALTERNATIV DEFENSE PLAN FOR WESTERN EUROPE".
PEACE RESEARCH SERIES NO. 3
Af: Bent Sørensen
- 96/85 "ASPEKTER VED KRAFTVARMEFORSYNING".
Af: Bjarne Lillethorup.
Vejleder: Bent Sørensen.
- 97/85 "ON THE PHYSICS OF A.C. HOPPING CONDUCTIVITY".
Af: Jeppe C. Dyre.
- 98/85 "VALGMULIGHEDER I INFORMATIONSSALDEREN".
Af: Bent Sørensen.
- 99/85 "Der er langt fra Q til R".
Projektrapport af: Niels Jørgensen og Mikael Klintorp.
Vejleder: Stig Andur Pedersen.
- 100/85 "TALSYSTEMETS OPBYGNING".
Af: Mogens Niss.
- 101/85 "EXTENDED MOMENTUM THEORY FOR WINDMILLS IN PERTURBATIVE FORM".
Af: Ganesh Sengupta.
- 102/85 OPSTILLING OG ANALYSE AF MATEMATISKE MODELLER, BELYST VED MODELLER OVER KØRS FODEROPTAGELSE OG - OMSÆTNING".
Projektrapport af: Lis Eilertzen, Kirsten Habekost, Lill Røn og Susanne Stender.
Vejleder: Klaus Grünbaum.

- 103/85 "ØDSLE KOLDKRIGERE OG VIDENSKABENS LYSE IDEER".
Projektrapport af: Niels Ole Dam og Kurt Jensen.
Vejleder: Bent Sørensen.
- 104/85 "ANALOGREGNEMASKINEN OG LORENZLIGNINGER".
Af: Jens Jøger.
- 105/85 "THE FREQUENCY DEPENDENCE OF THE SPECIFIC HEAT AT THE GLASS REANSITION".
Af: Tage Christensen.
- "A SIMPLE MODEL OF AC HOPPING CONDUCTIVITY".
Af: Jeppe C. Dyre.
Contributions to the Third International Conference on the Structure of Non - Crystalline Materials held in Grenoble July 1985.
- 106/85 "QUANTUM THEORY OF EXTENDED PARTICLES".
Af: Bent Sørensen.
- 107/85 "EN MYG GØR INGEN EPIDEMI".
- flodblindhed som eksempel på matematisk modellering af et epidemiologisk problem.
Projektrapport af: Per Hedegård Andersen, Lars Boye, Carsten Holst Jensen, Else Marie Pedersen og Erling Møller Pedersen.
Vejleder: Jesper Larsen.
- 108/85 "APPLICATIONS AND MODELLING IN THE MATHEMATICS CURRICULUM" - state and trends -
Af: Mogens Niss.
- 109/85 "COX I STUDIETIDEN" - Cox's regressionsmodel anvendt på studenteroplysninger fra RUC.
Projektrapport af: Mikael Wennerberg Johansen, Poul Katler og Torben J. Andreassen.
Vejleder: Jørgen Larsen.
- 110/85 "PLANNING FOR SECURITY".
Af: Bent Sørensen
- 111/85 "JORDEN RUNDT PÅ FLADE KORT".
Projektrapport af: Birgit Andresen, Beatriz Quinones og Jimmy Staal.
Vejleder: Mogens Niss.
- 112/85 "VIDENSKABELIGGØRELSE AF DANSK TEKNOLOGISK INNOVATION FREM TIL 1950 - BELYST VED EKSEMPLER".
Projektrapport af: Erik Odgaard Gade, Hans Hedal, Frank C. Ludvigsen, Annette Post Nielsen og Finn Physant.
Vejleder: Claus Bryld og Bent C. Jørgensen.
- 113/85 "DESUSPENSION OF SPLITTING ELLIPTIC SYMBOLS 11".
Af: Bernhelm Booss og Krzysztof Wojciechowski.
- 114/85 "ANVENDELSE AF GRAFISKE METODER TIL ANALYSE AF KONTINGENSTABELLER".
Projektrapport af: Lone Billmann, Ole R. Jensen og Anne-Lise von Moos.
Vejleder: Jørgen Larsen.
- 115/85 "MATEMATIKKENS UDVIKLING OP TIL RENESSANCEN".
Af: Mogens Niss.
- 116/85 "A PHENOMENOLOGICAL MODEL FOR THE MEYER-NELDEL RULE".
Af: Jeppe C. Dyre.
- 117/85 "KRAFT & FJERNVARMEOPTIMERING".
Af: Jacob Mørch Pedersen.
Vejleder: Bent Sørensen
- 118/85 "TILFÆLDIGHEDEN OG NØDVENDIGHEDEN IFØLGE PEIRCE OG FYSIKKEN".
Af: Peder Voetmann Christiansen
- 120/86 "ET ANTAL STATISTISKE STANDARDMODELLER".
Af: Jørgen Larsen
- 121/86 "SIMULATION I KONTINUERT TID".
Af: Peder Voetmann Christiansen.
- 122/86 "ON THE MECHANISM OF GLASS IONIC CONDUCTIVITY".
Af: Jeppe C. Dyre.
- 123/86 "GYMNASIEFYSIKKEN OG DEN STORE VERDEN".
Fysiklærerforeningen, IMFUA, RUC.
- 124/86 "OPGAVESAMLING I MATEMATIK".
Samtlige opgaver stillet i tiden 1974-jan. 1986.
- 125/86 "UVBY, 8 - systemet - en effektiv fotometrisk spektral-klassifikation af B-, A- og F-stjerner".
Projektrapport af: Birger Lundgren.
- 126/86 "OM UDVIKLINGEN AF DEN SPECIELLE RELATIVITETSTEORI".
Projektrapport af: Lise Odgaard & Linda Szkotak Jensen
Vejledere: Karin Beyer & Stig Andur Pedersen.
- 127/86 "GALOIS' BIDRAG TIL UDVIKLINGEN AF DEN ABSTRAKTE ALGEBRA".
Projektrapport af: Pernille Sand, Heine Larsen & Lars Frandsen.
Vejleder: Mogens Niss.
- 128/86 "SMÅRYB" - om ikke-standard analyse.
Projektrapport af: Niels Jørgensen & Mikael Klintorp.
Vejleder: Jeppe Dyre.
- 129/86 "PHYSICS IN SOCIETY"
Lecture Notes 1983 (1986)
Af: Bent Sørensen
- 130/86 "Studies in Wind Power"
Af: Bent Sørensen
- 131/86 "FYSIK OG SAMFUND" - Et integreret fysik/historie-projekt om naturanskuelsens historiske udvikling og dens samfundsmæssige betingethed.
Projektrapport af: Jakob Heckscher, Søren Brønd, Andy Wierød.
Vejledere: Jens Høyrup, Jørgen Vogelius, Jens Højgaard Jensen.
- 132/86 "FYSIK OG DANNEELSE"
Projektrapport af: Søren Brønd, Andy Wierød.
Vejledere: Karin Beyer, Jørgen Vogelius.
- 133/86 "CHERNOBYL ACCIDENT: ASSESSING THE DATA. ENERGY SERIES NO. 15.
Af: Bent Sørensen.
-
- 134/87 "THE D.C. AND THE A.C. ELECTRICAL TRANSPORT IN AsSeTe SYSTEM"
Authors: M.B.El-Den, N.B.Olsen, Ib Høst Pedersen, Petr Visčor
- 135/87 "INTUITIONISTISK MATEMATIKS METODER OG ERKENDELSESTEORETISKE FORUDSÆTNINGER"
MATEMATIKSPECIALE: Claus Larsen
Vejledere: Anton Jensen og Stig Andur Pedersen
- 136/87 "Mystisk og naturlig filosofi: En skitse af kristendommens første og andet møde med græsk filosofi"
Projektrapport af Frank Colding Ludvigsen
Vejledere: Historie: Ib Thiersen
Fysik: Jens Højgaard Jensen
- 137/87 "HOPMODELLER FOR ELEKTRISK LEDNING I UORDNEDE FASTE STOFFER" - Resume af licentiatafhandling
Af: Jeppe Dyre
Vejledere: Niels Boye Olsen og Peder Voetmann Christiansen.
- 119/86 "DET ER GANSKE VIST - - EUKLIDS FEMTE POSTULAT KUNNE NOK SKABE RØRE I ANDEDAMMEN".
Af: Iben Maj Christiansen
Vejleder: Mogens Niss.